

Ordering Physician:

John Doe, MD

1234 Main St. Anywhere, GA 30096 Accession #: Order #:

Reference #: Patient:

Date of Birth:

Reprinted: Comment:

Age:

Sex:

48

Female

A1012080012 G1234567

02/05/1962

07/09/2013

Sample Report

Date Collected: Date Received:

Egg, Whole

12/07/2010 12/08/2010

Date of Report: 12/08/2010

7704464583 Telephone:

Fax: 7704412237

0076 Allergix® Bloodspot IgG4 Food Antibodies 30 Profile

Methodology: ELISA

Almond

Mustard

Strawberry

	Foods to Avoid		
Negative	Mild	Moderate	Severe
	+1 and +2	+3 and +4	+5

Milk

Peanut

Aspergillus

Beef

Cantaloupe

Cashew

Chicken

Corn

Crab Garlic

Lobster

Oat

Orange

Pea, Green

Pinto Bean

Pork

Rice

Salmon

Shrimp

Soybean

Sunflower

Tomato Tuna

Turkey

Walnut

Wheat

Responses reflect IgG levels measured by ELISA with standardized food extracts. The assay yields semi-quantitative antibody concentrations for each food. The concentration readings are categorized into four reaction levels (Negative, Mild, Moderate, or Severe) corresponding to semi quantitative responses (0/1, +1, +2, +4 and +5) based on relative absorbance readings. The likelihood of adverse reactions to a given food increases as the response level for that food becomes more positive.

Florida Clinical Lab Lic. #800008124